

Non-equilibrium Phase-Ordering with a Global Conservation Law

A. D. Rutenberg*

Theoretical Physics, University of Oxford, Oxford OX1 3NP, United Kingdom

(February 1, 2008)

In all dimensions, infinite-range Kawasaki spin exchange in a quenched Ising model leads to an asymptotic length-scale $L \sim (\rho t)^{1/2} \sim t^{1/3}$ at $T = 0$ because the kinetic coefficient is renormalized by the broken-bond density, $\rho \sim L^{-1}$. For $T > 0$, activated kinetics recovers the standard asymptotic growth-law, $L \sim t^{1/2}$. However, at all temperatures, infinite-range energy-transport is allowed by the spin-exchange dynamics. A better implementation of global conservation, the microcanonical Creutz algorithm, is well behaved and exhibits the standard non-conserved growth law, $L \sim t^{1/2}$, at all temperatures.

Globally conserved dynamics offers a useful test of our theoretical understanding of phase-ordering [1] because it allows access to off-critical quenches without requiring diffusive “model-B” transport of the order-parameter. RG [2] arguments show that adding global conservation changes neither the critical dynamics nor the asymptotic growth law after a quench into the ordered phase. This has been confirmed by numerical studies [3]. The only change to non-conserved dynamics is to impose $\partial_t \langle \phi \rangle = 0$ where, using spin language, ϕ is the local magnetization ($\langle \phi \rangle$ is its spatial average). The standard “model-A” dissipative dynamics are $\partial_t \phi = -\Gamma \delta F / \delta \phi$, where $F[\{\phi\}] = \int d^d r [(\nabla \phi)^2 + (\phi^2 - 1)^2]$. The global constraint is imposed with a uniform magnetic field term, $H \langle \phi \rangle$, added to F . The field $H(t)$, effectively a lagrange multiplier, is decreased smoothly while the ordering progresses so as to maintain $\langle \phi \rangle$ constant.

With a global conservation law, $\langle \phi \rangle$ is a tunable parameter that does not change the growth-law but does affect scaled correlation functions and other aspects of the system. One can use this to explore what “universality” entails in non-equilibrium phase-ordering systems. For example, with global conservation in Ising models, Sire and Majumdar [4] have shown that the autocorrelation exponent λ depends on the net magnetization. This author [5] has found spatial anisotropy in scalar phase-ordering systems that depends strongly on $\langle \phi \rangle$.

However, there has been uncertainty about how to implement global conservation. Tuning a magnetic field to maintain $\langle \phi \rangle$ is impractical in computer simulations, due to the stochastic nature of most algorithms. Two alternative algorithms have been used for Ising models: a Creutz demon [6], and infinite-range Kawasaki spin-exchange [3,4,7]. Differences between these two implementations at low temperatures has led to some confusion in the past [4,7,8]. In this report we clarify the situation.

For a 2d Ising model on a square lattice, consider infinite-range Kawasaki exchange dynamics [3,4,7]: two randomly selected spins are exchanged under heat-bath dynamics, i.e. with probability $1/[1 + \exp(\Delta E/k_B T)]$, where ΔE is the energy change under the spin exchange. The exchange satisfies global conservation. However at low temperature the asymptotic dynamics are *acti-*

vated, and cross-over to a different growth law at zero-temperature.

At $T = 0$, no exchanges that increase the energy are accepted. Starting from random initial conditions, aggregation of spins takes place and the broken-bond density $\rho(t)$ will be proportional to the domain wall density at late times, $\rho \sim L^{-1}$. Isolated spins with all bonds broken, of number-density N , will aggregate rapidly onto existing domains, $\partial_t N \propto -\rho N$, and may be ignored. Spins which are part of domain walls can exchange, but only with the fraction ρ of the system that has broken bonds (i.e. that are also on domain walls). This rescales the effective kinetic coefficient by a factor of $\rho \sim L^{-1}$. For $T = 0$ infinite-range Kawasaki exchange we thus expect $L \sim (\rho t)^{1/2} \sim (t/L)^{1/2} \sim t^{1/3}$, where $t^{1/2}$ is the standard non-conserved growth [1]. This applies in any dimension.

For $T > 0$, thermal fluctuations provide partners for spin exchange and also allow exchange with bulk spins. These activated processes will dominate after the broken-bond density becomes comparable to the equilibrium average. After a quench, this will result in a crossover from intermediate $L \sim t^{1/3}$ growth to asymptotic $L \sim t^{1/2}$ growth. [The length-scale is extracted from the scaling of the spherically-averaged correlation function $C(r, t) = \langle \phi(x) \phi(x+r) \rangle = M^2 f(r/L)$, where M is the equilibrium bulk magnetization. L is chosen so that $f(1) = 1/2$.] Results for quenches to T_c will be similarly affected — the dynamical exponent will be underestimated while ρ relaxes towards equilibrium [9].

A renormalized kinetic coefficient alone will not affect *scaled* correlations. However Kawasaki exchange also allows infinite-range energy-transfer through the system. For example, at $T = 0$, an isolated spin (four broken bonds) may hop freely through the system by spin exchange with bulk spins (no broken bonds). Similarly, a “bump” on a flat interface (with one broken bond) may hop to any other flat interface of the system. Effects of this energy transport might show up at low-temperatures in non-universal effects such as spatial anisotropy in the late-stages of the quench [5]. It is unclear if this would affect the autocorrelation results of Sire and Majumdar [4], who used this algorithm at $T = 0$.

A faster and safer algorithm was proposed by Creutz

[6]. The system is coupled to a small spin reservoir (here of size 2). Single spin flips accepted by normal kinetic Ising model dynamics are subject to the additional condition that the required spin change (± 2) can be extracted from the reservoir. The dynamics of the system plus reservoir are microcanonical in the magnetization, and there are none of the questions of energy-transport that we have just raised for infinite-range Kawasaki exchange [10]. As a result the RG results directly apply and a growth-law of $L \sim t^{1/2}$ is indeed seen [3].

To check these arguments we simulated critical quenches, with $\langle \phi \rangle = 0$, for non-conserved and also for globally conserved systems with both Creutz and Kawasaki dynamics. For off-critical quenches similar results apply. In Fig. 1 we see the expected $t^{1/3}$ growth for $T = 0$ Kawasaki exchange, and reconfirm $t^{1/2}$ growth for the other cases. To test the rescaling of the kinetic prefactor in the Kawasaki algorithm, we show a quench to $T = 0.9T_c$ and also $L/\sqrt{\rho}$ for the quench to $T = 0$. Both of these exhibit the expected $t^{1/2}$ growth. In Fig. 2, we see that all of the models have similar spherically-averaged scaled correlations. [Anisotropies are too small to see for these critical quenches [5].] The correlations for the Kawasaki model are slightly different, however this seems to be a transient effect. [Strong finite-size effects in the $T = 0.9T_c$ quench, after the latest time shown, limit the duration of the simulation.]

In conclusion, we see that an infinite-range Kawasaki exchange implementation of global conservation laws has activated kinetics, with an anomalous $L \sim t^{1/3}$ growth at $T = 0$. In addition, Kawasaki dynamics leads to long-range energy transport. The Creutz algorithm, on the other hand, is not activated and has no long-range energy transport. It provides a faster and better-controlled implementation of global conservation laws, and should exhibit the expected $t^{1/2}$ growth at all temperatures.

This work was supported by EPSRC grant GR/J78044.

Rutenberg, *ibid* **76**, 158 (1996).

- [9] This may be the cause of the anomalously small dynamical critical exponent seen by Tamayo and Klein [7] with infinite-range Kawasaki exchange.
- [10] The Creutz algorithm must not be implemented with a sublattice update since the sweep direction generates anomalous spatial anisotropies.




FIG. 1. Length-scale $L(t)$ for non-conserved (stars), Creutz (plus), Kawasaki $T = 0.9T_c$ (triangle), and Kawasaki $T = 0$ (circle). The system sizes are 2048^2 , 1024^2 , 512^2 , and 512^2 , respectively, with at least 20 samples in each system. We also show $L/\sqrt{\rho}$ for the $T = 0$ Kawasaki quench (squares), which goes as $t^{1/2}$, as expected. The upper and lower lines show $t^{1/2}$ and $t^{1/3}$, respectively.




FIG. 2. Scaled correlations, divided by the equilibrium magnetization squared, from the latest times of simulations shown in the previous figure. Solid, dashed, dot-dashed, and dotted lines are non-conserved, Creutz $T = 0$, Kawasaki $T = 0$, and Kawasaki $T = 0.9T_c$, respectively.

* adr@thphys.ox.ac.uk

- [1] For a review, see A. J. Bray, *Adv. Phys.* **43**, 357 (1994).
- [2] A. J. Bray, *Phys. Rev. Lett.* **62**, 2841 (1989); *Phys. Rev. B* **41**, 6724 (1990). See also [1].
- [3] J. F. Annett and J. R. Banavar, *Phys. Rev. Lett.* **68**, 2941 (1992); L. L. Moseley *et al.*, *J. Stat. Phys.* **67**, 813 (1992).
- [4] C. Sire and S. N. Majumdar, *Phys. Rev. E* **52**, 244 (1995).
- [5] A. D. Rutenberg, cond-mat/9602097.
- [6] M. Creutz, *Phys. Rev. Lett.* **50**, 1411 (1983).
- [7] P. Tamayo and W. Klein, *Phys. Rev. Lett.* **63**, 2757 (1989); *ibid.* **66**, 2049 (1991); A. J. Bray, *ibid.* **66**, 2048 (1991).
- [8] S. B. Goryachev, *Phys. Rev. Lett.* **72**, 1850 (1994); A. D.